



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Br

Search PubMed for Go Clear Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Enzyme Microb Technol. 1993 Nov;15(11):950-8.

Related Articles, Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

**A novel NADH-dependent carbonyl reductase with an extremely broad substrate range from Candida parapsilosis: purification and characterization.****Peters J, Minuth T, Kula MR.**

Institut für Enzymtechnologie, Heinrich-Heine-Universität Düsseldorf, Jülich, Germany.

SDS 32 kDa  
gel 85 kDa  
64 16  
170 43

A novel oxidoreductase catalyzing the NADH-dependent reduction of a variety of carbonyl compounds, especially keto esters, was found in *Candida parapsilosis* DSM 70125. The enzyme was purified by fractional poly (ethylene glycol) precipitation, anion exchange, and affinity chromatography. The enzyme was enriched about 3100-fold and appeared to be homogeneous as judged by native and sodium dodecyl sulfate gel electrophoresis. The carbonyl reductase from *C. parapsilosis* is a dimeric enzyme with an apparent molecular mass of about 135 kDa. Important properties concerning the application of the enzyme are the relatively broad pH optimum between pH 6.5 and 9.0, temperature optimum between 36 and 42 degrees C, and good stability. Besides keto esters, the new enzyme reduces other aliphatic, aromatic, and cyclic ketones, as well as aldehydes and ketoacetals with high reaction rates. 4-Halo-3-hydroxybutanoates, which are promising chiral intermediates for the chemical synthesis of L-carnitine, alkaloids and pharmaceuticals, are now accessible by enzymatic reduction, as well as several phenyl-ethanol derivatives, which are important for the synthesis of pharmaceuticals and agrochemicals. The preparative applicability of the enzyme was demonstrated in a coupled enzyme system with regeneration of coenzyme. Methyl 3-oxobutanoate was converted into methyl (S)-(+)-3-hydroxybutanoate (98.5% ee), a versatile chiral building block for the synthesis of pheromones and different antibiotics.

PMID: 7764255 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Aug 30 2004 06:52:01

h      cb      hg   e   e   e   fcg   c      e   e   e   b   b   e      b   c